

NON-DIAGONAL PARTON DISTRIBUTION FUNCTIONS¹

M. DIEHL

DAPNIA/SPhN, CEA/Saclay, 91191 Gif sur Yvette CEDEX, France
E-mail: mdiehl@cea.fr

and

T. GOUSSET

SUBATECH², B.P. 20722, 44307 Nantes CEDEX 3, France
E-mail: gousset@subatech.in2p3.fr

Abstract

We show why the time ordering in the definition of non-diagonal parton distributions can be omitted and discuss the physics implications.

1 The physics context

The idea that the gluon distribution in the proton is relevant to diffractive photon-proton reactions [1] has been very fruitful and now been applied to a variety of processes: diffractive production of vector mesons, of a real photon or Z , of dijets and of open charm. It can be generalised to the non-diffractive region, where gluon exchange is no longer dominant and the quark distribution is probed [2, 3]. Then one can also exchange quantum numbers and consider final states such as $p\pi^0$ or $n\pi^+$. In all cases one exchanges a quark-antiquark or a gluon pair, attached to a parton distribution S on the proton and to a hard scattering H on the photon side as shown in Fig. 1; it is required that the photon virtuality or the final state provide a hard scale while t is small. For certain processes this factorisation has been derived in QCD [3].

To make the transition from γ^* to X possible the hadron momenta on either side of the blob S must be different, in contrast to usual parton distributions. We thus have new

¹Talk given at the International Workshop on Deep Inelastic Scattering and QCD (DIS 98), Brussels, Belgium

²Unité Mixte de l'Université de Nantes, de l'Ecole des Mines de Nantes et du CNRS




Figure 1: The factorisation of $\gamma^* + p \rightarrow X + p$ into a non-diagonal parton distribution S and a hard scattering H . If X is a meson, H includes a distribution amplitude describing the formation of the meson from its valence quarks. The partons have momenta k and k' with plus-components $k^+ = x_1 p^+$ and $k'^+ = x_2 p^+$.

quantities, non-diagonal parton distributions, which depend on t and on the momentum fractions x_1 and x_2 of the partons. Their difference $x_1 - x_2$ is fixed to be $\xi = [(p-p') \cdot q] / [p \cdot q]$ by kinematics.

2 Time ordering: why we want to drop it ...

Taking for X in Fig. 1 the incoming γ^* and cutting in the s -channel we obtain the diagram for the inclusive $\gamma^* p$ cross section via the optical theorem. The blob S , described now by usual parton densities, is then cut, whereas the non-diagonal parton distributions appear in the diagram of an *amplitude* where S is not cut. On a formal level ordinary quark or gluon distributions are Fourier transformed matrix elements of products $\bar{\psi}_\beta(0) \psi_\alpha(z) |_{z^2=0}$ or $G_{\mu\nu}(0) G_{\rho\sigma}(z) |_{z^2=0}$ of field operators, whereas the non-diagonal ones appear with *time ordered* products.

It can however be shown that, because the separation z of the fields is lightlike, one can leave out the time ordering, in other words it is the same whether one cuts the blob S or not [4]. Before giving the idea of the proof let us point out why this property is so important:

- As non-diagonal distributions can be expressed without time ordering they become equal to the usual parton distributions in the limit $p = p'$.
- If the blob S is cut one can show that, apart from convention dependent phases, the corresponding parton distributions are *real valued* due to time reversal invariance.
- Since the blob S is “already cut” the real and imaginary parts of the $\gamma^* p$ amplitude in Fig. 1 are obtained just from the real and imaginary parts of the hard scattering H , convoluted with the same soft quantity. This is needed to obtain a simple dispersion relation for the amplitude [3].
- While the momentum fractions x_1, x_2 can a priori range from $-\infty$ to $+\infty$ in the loop integral of Fig. 1, they are restricted to a finite interval because the lower blob is effectively cut. This is crucial for a parton interpretation: no parton in a hadron can be faster than the hadron itself.
- An ordinary parton distribution can be interpreted as the squared amplitude for the

emission of a specified parton, summed over all configurations of spectator partons, which go across the cut of the blob S . Since non-diagonal distributions are cut as well they have a similar interpretation. Because x_1 and x_2 are different they involve however the *interference* of two different amplitudes. In this sense they contain essentially new information on the proton structure.

The question of time ordering is also relevant for higher twist parton distributions, *i.e.* soft blobs with more than two parton lines. For the diagonal case, $p = p'$, Jaffe has shown that time ordering can be dropped in leading and nonleading twist distributions [5].

3 ... and why we are allowed to

The idea of our proof is to use the analytic properties of scattering amplitudes, following Landshoff and Polkinghorne [6]. Let us work in the collision c.m. with the 3-axis along p and introduce for any vector v its light cone coordinates $v^\pm = (v^0 \pm v^3)/\sqrt{2}$ and its transverse part $v_T = (v^1, v^2)$. The parton distributions can be expressed in terms of

$$\int dk^- d^2k_T \mathcal{A}|_{k^+=x_1 p^+} \quad , \quad (1)$$

where $\mathcal{A}$ is the proton-parton scattering amplitude depicted by the blob S , given by $\mathcal{A} = \int d^4z e^{ik \cdot z} \langle p' | T \bar{\psi}_\beta(0) \psi_\alpha(z) | p \rangle$ if for definiteness we take quark distributions. In position space the integral over k^- corresponds to setting $z^+ = 0$ in the matrix element, and the combined integral over k^- and k_T puts z on the light cone. It turns out that the integration over k^- is enough for our purpose (we may thus also consider k_T -unintegrated distributions).

According to the usual ideas on analyticity the amplitude $\mathcal{A}$ has singularities (poles and branch cuts) in the parton virtualities k^2 , k'^2 and in the Mandelstam variables $s = (-k+p)^2$, $u = (k+p')^2$; in the standard convention they lie slightly below the real axis of the variable concerned. Expressing each variable in terms of the components of p , p' and k we can map these singularities onto the complex k^- -plane. We find that their location is controlled by the fractions x_1 and $x_2 = x_1 - \xi$ and have several cases:

- If $x_1 > 1$ or $x_1 < \xi - 1$ then all singularities lie on the same side of the real k^- -axis. Thus we can close the integration contour by an infinite semicircle in the other half-plane without encircling any singularity, and obtain a zero integral. Under the assumption that $\mathcal{A}$ vanishes fast enough as $|k^-| \rightarrow \infty$ at fixed k^+ and k_T the semicircle does not contribute and the integral along the real k^- -axis is zero itself. We thus find that our distributions are only nonzero in the interval $\xi - 1 < x_1 < 1$.
- For $\xi < x_1 < 1$ we find the singularities in s above and all others below the real k^- -axis. Closing the contour in the upper half-plane our integral is given by the discontinuity of $\mathcal{A}$ across the s -cut, which according to the Cutkosky rules corresponds to cutting the soft amplitude in the s -channel, as shown in Fig. 2 (a). This cut amplitude can be written in terms of $\langle p' | \bar{\psi}_\beta(0) | I \rangle \langle I | \psi_\alpha(z) | p \rangle$ summed over all intermediate states I ; using the closure property we find just the product $\bar{\psi}_\beta(0) \psi_\alpha(z)$ without time ordering as we wanted.




Figure 2: Different ways to cut the proton-parton amplitude $\mathcal{A}$: (a) in the s -channel, (b) in the parton virtuality k'^2 . I and I' denote the intermediate states.

- For $\xi - 1 < x_1 < 0$ we can pick up in a similar way the u -channel cut of the soft amplitude and obtain a matrix element of $-\psi_\alpha(z)\bar{\psi}_\beta(0)$. While for $\xi < x_1 < 1$ we had the emission of a quark with momentum fraction x_1 and its reabsorption with fraction x_2 we now have emission of an antiquark with fraction $-x_2$ and its reabsorption with fraction $-x_1$.
- The region $0 < x_1 < \xi$ has the particularity that x_1 is positive but x_2 negative; in a parton picture one has now the transition from a proton p to a proton p' and a quark-antiquark pair. In the k^- -plane the singularities in s and k'^2 lie above, those in u and k^2 below the real axis. If we chose to pick up the former two we must cut the soft amplitude in the s -channel as before, but we must also cut in k'^2 , as shown in Fig. 2 (b). Altogether we find precisely the sum over intermediate states I and I' needed to go from a time ordered product of quark fields to an ordinary one.

In summary, we can show under fairly general assumptions on the amplitude $\mathcal{A}$ that in parton distributions the time ordering of fields is irrelevant because the component z^+ of their separation is zero, in other words because parton distributions are integrated over the minus component of the parton momentum.

References

- [1] J. Bartels and M. Loewe, *Z. Phys. C* **12**, 263 (1982); M.G. Ryskin, *Z. Phys. C* **57**, 89 (1993); S.J. Brodsky *et al*, *Phys. Rev. D* **50**, 3134 (1994).
- [2] X. Ji, *Phys. Rev. Lett.* **78**, 610 (1997); *Phys. Rev. D* **55**, 7114 (1997); A.V. Radyushkin, *Phys. Rev. D* **56**, 5524 (1997).
- [3] J.C. Collins, L. Frankfurt and M. Strikman, *Phys. Rev. D* **56**, 2982 (1997); J.C. Collins and A. Freund, hep-ph/9801262.
- [4] M. Diehl and T. Gousset, hep-ph/9801233, to appear in *Phys. Lett. B*.
- [5] R.L. Jaffe, *Nucl. Phys. B* **229**, 205 (1983).
- [6] P.V. Landshoff and J.C. Polkinghorne, *Phys. Rep.* **5**, 1 (1972).